

Author and Reference	Year of publication	Number of patients	Previous (failed) treatments	Duration (days)	Eradication (%)
Gatta et al. [275]	2005	151	1 st : PPI+C+A or M 2 nd : PPI+C+A or M;	7-14	76
Gisbert et al [276]	2006	100	1 st : PPI+C+A 2 nd : Q	10	60
Gisbert et al [277]	2006	20	1 st : PPI+C+A 2 nd : Q	10	85
Hsu et al [262]	2008	37	1 st : PPI+C+A 2 nd : Q	10	84
Rokkas et al [278]	2009	30	1 st : PPI+C+A 2 nd : Q	10	70
Gisbert et al [279]	2012	200	1 st : PPI+C+A 2 nd : Q	10	68
Burgos-Santamaría [280]	2019	65	1 st : PPI+C+A 2 nd : Q	7-14	75
Burgos-Santamaría [280]	2019	19	1 st : PPI+C+A 2 nd : Q	14	84
Burgos-Santamaría [281]	2021	39	1 st : PPI+C+A±M 2 nd : Q	10-14	74
Burgos-Santamaría [281]	2021	28	1 st : PPI+C+A±M 2 nd : Q	14	75

PPI: proton pump inhibitor; C: clarithromycin; M: metronidazole; A: amoxicillin; Q: bismuth-containing quadruple therapy (PPI, bismuth, tetracycline and M).

[†]Bismuth was added to the triple regimen with a proton pump inhibitor, amoxicillin, and levofloxacin.